

## Part II Algebraic geometry

### Example Sheet I, 2015

In all problems, you may assume that we are working over an algebraically closed field  $k$ .

1. Let  $f : X \rightarrow Y$  be a morphism of affine schemes. Show that  $f$  is a continuous map in the Zariski topology.

2. Let  $Y \subseteq \mathbf{A}^2$  be the curve given by  $xy = 1$ . Show that  $Y$  is not isomorphic to  $\mathbf{A}^1$ . Find all morphisms  $\mathbf{A}^1 \rightarrow Y$  and  $Y \rightarrow \mathbf{A}^1$ .

3. Let  $Y \subseteq \mathbf{A}^3$  be the set  $\{(t, t^2, t^3) \mid t \in k\}$ . Show that  $Y$  is an affine variety, determine  $I(Y)$ , and show that  $A(Y)$  is a polynomial ring in one variable.  $Y$  is called the *twisted cubic*.

4. Let  $Y = Z(x^2 - yz, xz - x)$ . Show that  $Y$  has 3 irreducible components. Describe them, and their corresponding prime ideals.

5. Show that any non-empty open subset of an irreducible variety is dense. Show that if an affine variety is Hausdorff, it consists of a single point.

6. A topological space is called *Noetherian* if it satisfies the descending chain condition for closed subsets. Show that affine varieties are Noetherian in the Zariski topology.

7. Show that if  $X \subseteq \mathbf{A}^n$ ,  $Y \subseteq \mathbf{A}^m$  are affine varieties, then  $X \times Y \subseteq \mathbf{A}^n \times \mathbf{A}^m = \mathbf{A}^{n+m}$  is a Zariski closed subset of  $\mathbf{A}^{n+m}$ , by explicitly writing  $I(X \times Y)$  in terms of  $I(X) = (f_1(x_1, \dots, x_n), \dots, f_t(x_1, \dots, x_n))$  and  $I(Y) = (h_1(y_1, \dots, y_m), \dots, h_s(y_1, \dots, y_m))$ .

8. Let  $Y \subseteq \mathbf{A}^3$  be the set  $\{(t^3, t^4, t^5) \mid t \in k\}$ . Show that  $Y$  is an affine variety, and determine  $I(Y)$ . Show  $I(Y)$  cannot be generated by two elements.

9. Show that there are no non-constant morphisms from  $\mathbf{A}^1$  to  $E = Z(y^2 - x^3 + x)$ .

10. Let  $f \in k[x_1, \dots, x_n]$  be an irreducible polynomial, and consider  $Y = Z(yf - 1) \subseteq \mathbf{A}^{n+1}$ , with coordinates  $x_1, \dots, x_n, y$ . Show that  $Y$  is irreducible. Show that the projection  $\mathbf{A}^{n+1} \rightarrow \mathbf{A}^n$  given by  $(x_1, \dots, x_n, y) \mapsto (x_1, \dots, x_n)$  induces a morphism  $Y \rightarrow \mathbf{A}^n$  which is a homeomorphism to its image  $D(f) := \{(a_1, \dots, a_n) \in \mathbf{A}^n \mid f(a_1, \dots, a_n) \neq 0\}$ . This gives the Zariski open set  $D(f)$  the structure of an algebraic variety.

11. Show that  $G = GL_n(k)$  is an affine variety, and that the multiplication and inverse maps are morphisms of algebraic varieties. We say  $G$  is an *affine algebraic group*. Show that if  $G$  is an affine algebraic group, and  $H$  is a subgroup which is also a closed subvariety of  $G$ , then  $H$  is also an affine algebraic group.

12. Let  $Mat_{n,m}$  be the set of  $n$  by  $m$  matrices with coefficients in  $k$ ; this set can be identified with  $\mathbf{A}^{nm}$  in the obvious way.

a) Show that the set of 2 by 3 matrices of rank  $\leq 1$  is an algebraic set.

b) Show that the matrices in  $Mat_{n,m}$  of rank  $\leq r$  is an algebraic set.

13. Let  $f, g \in k[x, y]$  be polynomials, and suppose  $f$  and  $g$  have no common factor. Show there exists  $u, v \in k[x, y]$  such that  $uf + vg$  is a non-zero polynomial in  $k[x]$ .

Now let  $f \in k[x, y]$  be irreducible. The variety  $Z(f)$  is called an affine *plane curve*. Show that any proper subvariety of  $Z(f)$  is finite.

14. Let  $A$  be a  $k$ -algebra. We say  $A$  is *finitely generated* if there is a surjective  $k$ -algebra homomorphism  $k[x_1, \dots, x_n] \rightarrow A$  for some  $n$ . Now suppose that  $A$  is a finitely generated  $k$ -algebra which is also an integral domain. Show that there is an affine variety  $Y$  with  $A$  isomorphic to  $A(Y)$  as  $k$ -algebras.

15. Let  $G = \mathbf{Z}/2\mathbf{Z}$  act on  $k[x, y]$  by sending  $x \mapsto -x, y \mapsto -y$ . Show that the algebra of invariants  $k[x, y]^G$  (the subring of polynomials left fixed by this action) defines an affine subvariety  $X$  of  $\mathbf{A}^3$  by explicitly computing this ring of invariants.  $X$  is called the *rational double point*.

What is the relation of the points of  $X$  to the orbits of  $G$  acting on  $\mathbf{A}^2$ ?